

Normal Newborn Admission

Order Set Directions:

 = Performance Indicator

(√)-Check orders to activate. Orders with a pre-checked box will be followed unless lined out.
Initial every order page & Sign, date, and time the last page.

Length _____ Weight _____

Nursing

- Transcutaneous Bilirubin prior to discharge (if greater than 7, do serum bilirubin)
- Newborn Hearing Screening after 12 hours of Age (after 2nd referral schedule O.P. hearing screening in one week post discharge)
- Bedside glucose if infant experiencing signs of hypoglycemia
- Pulse oximetry screening at 24 hours of age
- Follow standard nursing policy for Care of the Normal Newborn

Signs of hypoglycemia

- ◆ Jitteriness
- ◆ Cyanosis
- ◆ Apnea
- ◆ Hypothermia
- ◆ Poor body tone
- ◆ Poor feeding
- ◆ Lethargy
- ◆ Seizures

Medications

Analgesics:

sucrose (SWEET-EASE)

- 1 dose PO once 2 minutes prior to a painful procedure; use a pacifier dipped in a 24% solution; maximum 2ml

Antibacterial Prophylactic Agents: Ophthalmia Neonatorum

erythromycin 5 mg/g eye ointment (ILOTYCIN)

- 0.5 centimeter in each eye once upon admission, if not given in delivery room or operating room

Immunizations

hepatitis B virus vaccine (preserv free) (ENGERIX-B)

- 10 ucg intramuscularly once

Topical Agents

zinc oxide ointment (DESITIN)

- applied topically 3 times a day PRN for diaper rash

AQUAPHOR

- applied topically 3 times a day PRN for dry skin

white petrolatum ointment

- applied topically PRN with diaper changes after circumcision

Vitamin K Preparations

phytonadione (AQUAMEPHYTON)

- 0.5 mg intramuscularly once

Laboratory

- Direct Coombs if mother is type O Positive or no prenatal testing for antibodies performed
- Cord Blood Panel if mom is Rh Negative
- Metabolic Screening after 24 hours of age
- Rapid HIV with undocumented maternal HIV status ASAP after birth

Provider initials: _____



Normal Newborn Admission

Order Set Directions:

 = Performance Indicator

(√)-Check orders to activate. Orders with a pre-checked box will be followed unless lined out.
Initial every order page & Sign, date, and time the last page.

Other

Circumcision on _____ at _____

Analgesics: Topical

lidocaine-prilocaine 2.5% topical cream (EMLA)

- applied topically once 1 hour prior to circumcision; gestational age greater than or equal to 37 weeks; maximum total dose 1 gram; maximum application area 10 centimeters squared; maximum application time 1 hour

Other Orders

Provider Signature: _____ Date: _____ Time: _____

